Π

(Acts adopted under the EC Treaty/Euratom Treaty whose publication is not obligatory)

DECISIONS

COMMISSION

COMMISSION DECISION

of 13 April 2007

on the implementation of surveillance programmes for avian influenza in poultry and wild birds to be carried out in the Member States and amending Decision 2004/450/EC

(notified under document number C(2007) 1554)

(2007/268/EC)

THE COMMISSION OF THE EUROPEAN COMMUNITIES,

Having regard to the Treaty establishing the European Community,

Having regard to Council Decision 90/424/EEC of 26 June 1990 on expenditure in the veterinary field (1) and in particular the fourth subparagraph of Article 24(2) and Article (10) thereof,

Having regard to Council Directive 2005/94/EC of 20 December 2005 on Community measures for the control of avian influenza and repealing Directive 92/40/EEC (2) and in particular Article 4(2) thereof,

Whereas:

- Decision 90/424/EEC of 26 June 1990 on expenditure in (1)the veterinary field lays down the procedures governing the Community's financial contribution towards programmes for the eradication, control and monitoring of animal diseases.
- Decision 90/424/EEC, as amended by Decision (2)2006/53/EC (³), provides that Community financial assistance may be granted to Member States for those eradication measures carried out by the Member States to combat low pathogenic avian influenza (LPAI) strains known to have the potential to mutate to highly pathogenic avian influenza (HPAI). In addition, Article

24(2) of Decision 90/424/EEC provides that Member States are to submit to the Commission, each year by 30 April at the latest, the annual or multi-annual programmes starting in the following year for which they wish to receive a financial contribution from the Community.

- (3) Council Directive 92/40/EEC of 19 May 1992 introducing Community measures for the control of avian influenza (4) defines the Community control measures to be applied in the event of an outbreak of highly pathogenic avian influenza (HPAI) in poultry. However, it does not provide for the control of low pathogenic avian influenza (LPAI) of H5 and H7 subtypes or for regular surveillance of that disease in poultry and wild birds.
- Since 2002, Member States have been implementing (4)mandatory surveys for avian influenza in domestic poultry by submitting yearly surveillance programmes to the Commission, as provided for in Commission Decisions 2002/649/EC (⁵), 2004/111/EC (⁶), 2005/464/EC (7) and 2006/101/EC (8).
- Directive 2005/94/EC provides for certain preventive (5)measures relating to the surveillance and early detection of avian influenza. The final date for the transposition of that Directive by the Member States is 1 July 2007 and it provides for the repeal of Directive 92/40/EEC as of that date.

(8) OJ L 46, 16.2.2006, p. 40.

^{(&}lt;sup>1</sup>) OJ L 224, 18.8.1990, p. 19. Decision as last amended by Council Decision 2006/965/EC (OJ L 397, 30.12.2006, p. 22).

⁽²⁾ OJ L 10, 14.1.2006, p. 16.

⁽³⁾ OJ L 29, 2.2.2006, p. 37.

^{(&}lt;sup>4</sup>) OJ L 167, 22.6.1992, p. 1. Directive as last amended by Directive 2006/104/EC (OJ L 363, 20.12.2006, p. 352).

⁽⁵⁾ OJ L 213, 9.8.2002, p. 38.
(6) OJ L 32, 5.2.2004, p. 20. Decision as amended by Decision 2004/615/EC (OJ L 278, 27.8.2004, p. 59).

 ⁽⁷⁾ OJ L 164, 24.6.2005, p. 52. Decision as amended by Decision 2005/726/EC (OJ L 273, 19.10.2005, p. 21).

- (6) The Community measures for the control of avian influenza provided for in Directive 2005/94/EC also cover the control of outbreaks of LPAI caused by avian influenza of H5 and H7 subtypes in poultry. In order to detect the possible circulation of those viruses in poultry flocks, compulsory surveillance programmes are to be implemented by the Member States. Those control measures aim at preventing the spread of LPAI of H5 and H7 subtypes before they become widespread in the domestic poultry population so that the risk of a mutation into HPAI with possibly devastating consequences may be prevented.
- (7) Directive 2005/94/EC also provides for surveillance programmes to be carried out in wild birds in order to contribute, on the basis of a regularly updated risk assessment, to the knowledge on the threats posed by wild birds in relation to any influenza virus of avian origin in birds.
- (8) It is important to further strengthen surveillance activities given recent developments as regards the widespread occurrence of HPAI H5N1 in wild birds in Europe, taking into account the results of the surveys carried out in the Member States between 2003 and 2006 and the scientific work recently undertaken by European Food Safety Authority (EFSA) (¹) in collaboration with the ORNIS Scientific Working Group of the European Commission's Environment Directorate-General. Those bodies will continue their work and outcomes thereof may lead to further updating.
- (9) When implementing surveillance programmes in wild birds full regard shall be paid to the requirements of Council Directive 79/409/EEC (²) on the protection and conservation of all naturally occurring wild bird species in the Community.
- (10) Commission Decision 2004/450/EC of 29 April 2004 laying down standard requirements for the content of applications for Community financing for programmes for the eradication, monitoring and control of animal diseases (³), lays down standard requirements for the content of applications for Community financing programmes for the eradication, monitoring and control of animal diseases.
- (11) As Decision 90/424/EC, now provides that a Community financial contribution is to be granted for expenditure incurred by the Member States for the financing of

- (²) OJ L 103, 25.4.1979, p. 1. Directive as last amended by Directive 2006/105/EC (OJ L 363, 20.12.2006, p. 368).
 (³) OJ L 155, 30.4.2004, p. 90; corrected by OJ L 92, 12.4.2005, p. 16.
- (3) OJ L 155, 30.4.2004, p. 90; corrected by OJ L 92, 12.4.2005, p. 16. Decision as amended by Decision 2006/282/EC (OJ L 104, 13.4.2006, p. 40).

national programmes for the eradication, control and monitoring of certain animal diseases, including avian influenza, Member States may submit to the Commission surveillance programmes for avian influenza for a financial contribution from the Community not later than 30 April each year pursuant to Article 24(2) of Decision 90/424/EEC. Decision 2004/450/EC should be amended to lay down standard requirements for the content of applications for Community financing for avian influenza surveillance programmes.

- (12) Decision 2004/450/EC should therefore be amended accordingly.
- (13) The measures provided for in this Decision are in accordance with the opinion of the Standing Committee of the Food Chain and Animal Health,

HAS ADOPTED THIS DECISION:

Article 1

Approval of surveillance programmes for avian influenza

The surveillance programmes for avian influenza in poultry and wild birds to be carried out by Member States, in accordance with Article 4(1) of Directive 2005/94/EC, shall comply with the guidelines set out in Annexes I and II to this Decision.

Article 2

Amendments to Decision 2004/450/EC

Decision 2004/450/EC is amended as follows:

- 1. in Article 1, the following point (c) is added:
 - '(c) in respect of the animal disease referred to in Annex I, Part C, at least the information set out in Annex IV.';
- 2. in Annex I, the following Part C is added:

'PART C

Disease referred to in Article 1(c)

avian influenza.';

3. A new Annex IV, the text of which is set out in Annex III to the present Decision, is added.

^{(&}lt;sup>1</sup>) Scientific opinion on 'Migratory birds and their possible role in the spread of highly pathogenic avian influenza' (EFSA, 12 May 2006) and its addendum (11 December 2006).

Article 3

This Decision is addressed to the Member States.

Done at Brussels, 13 April 2007.

For the Commission Markos KYPRIANOU Member of the Commission

ANNEX I

Guidelines on the implementation of surveillance programmes for avian influenza in poultry to be carried out in the Member States

A. Objectives, general requirements and criteria for surveillance

A.1. Objectives

Serological surveillance for LPAI subtypes H5 and H7 in poultry shall aim at:

- 1. Detecting sub-clinical infections with LPAI of subtypes H5 and H7 thereby complementing early detection systems and subsequently preventing possible mutation of these viruses to HPAI.
- 2. Detecting infections of LPAI H5 and H7 subtypes in specifically targeted poultry populations at specific risk for infection due to their husbandry system or the susceptibility of specific species.
- 3. Contributing to the demonstration of a free status of a certain country, region or compartment from notifiable avian influenza in the frame of international trade according to OIE rules.
- A.2. General requirements and criteria
- 1. Sampling shall not extend beyond 31 December of the year of implementation of the programme. For poultry, sampling shall cover a period appropriate to production periods for each poultry category as required.
- 2. In order to save resources, samples collected for other purposes are recommended.
- 3. Testing of samples shall be carried out at national laboratories for avian influenza (NL) in Member States or by other laboratories authorised by the competent authorities and under the control of the NL.
- 4. All results (both serological and virological) shall be sent to the Community Reference Laboratory for Avian Influenza (CRL) for collation. A good flow of information must be ensured. The CRL shall provide technical support and keep an enlarged stock of diagnostic reagents.
- 5. All avian influenza virus isolates shall be submitted to the CRL in accordance with Community legislation, unless a derogation according to paragraph 4 of Chapter V (Differential diagnosis) in the diagnostic manual laid down in Decision 2006/437/EC (¹) is granted. Viruses of H5/H7 subtype shall be submitted without delay and shall be subjected to the standard characterisation tests (nucleotide sequencing/IVPI) according to that Diagnostic Manual.
- 6. Whenever possible, NLs shall submit to the CRL, H5 or H7 positive sera collected from Anseriformes in order that an archive be established to facilitate future test development.

B. Surveillance for avian influenza in poultry

- 1. All positive findings shall be retrospectively investigated at the holding and the conclusions of this investigation shall be reported to the Commission and the CRL.
- 2. Specific protocols to accompany the sending of material to the CRL and reporting tables for collection of surveillance data shall be provided by the CRL. In those tables the laboratory testing methods used shall be indicated. The tables provided shall be used to submit results in a single document.

- 3. Blood samples for serological examination shall be collected from all species of poultry including those reared in free-range systems, from at least 5 to 10 birds (except ducks geese and quail) per holding, and from the different sheds, if more than one shed is present on a holding. In case of several sheds the sample size per holding should be increased appropriately. It is recommended to take at least five birds per shed.
- 4. Sampling shall be stratified throughout the territory of the whole Member State, so that samples can be considered as representative for the whole of the Member State, taking into account:
 - (a) the number of holdings to be sampled (excluding ducks, geese and turkeys); that number shall be defined so as to ensure the identification of at least one infected holding if the prevalence of infected holdings is at least 5 %, with a 95 % confidence interval (see Table 1); and
 - (b) the number of birds sampled from each holding shall be defined so as to ensure 95 % probability of identifying at least one positive bird if the prevalence of sero-positive birds is ≥ 30 %.
- 5. Based on a risk assessment and the specific situation in the Member State concerned, the sampling design shall also consider:
 - (a) the types of production and their specific risks, shall be targeted to free range production, outdoor keeping and backyard flocks plus taking into account other factors such as multi-age, use of surface water, a relatively longer life span, the presence of more than one species on the holding or other relevant factors;
 - (b) the number of turkey, duck and goose holdings to be sampled shall be defined to ensure the identification of at least one infected holding if the prevalence of infected holdings is at least 5 %, with a 99 % confidence interval (see Table 2);
 - (c) where significant number of holdings producing game, ratites and quails are present in a Member State they shall be included in the programme. With regard to quails only adult (or laying) breeders shall be sampled;
 - (d) the time period for sampling shall coincide with seasonal production. However, where appropriate, sampling can be adapted to other identified periods at local level, during which time the presence of other poultry hosts on a holding might pose a greater risk for disease introduction;
 - (e) in case a significant number of backyard flocks are present, surveillance could be extended to them;
 - (f) Member States that must carry out sampling for Newcastle disease to maintain their status as Newcastle disease non-vaccinating countries in accordance with Commission Decision 94/327/EC (¹) may utilise these samples from breeding flocks for the surveillance of H5/H7 antibodies.

Τ	`ab	le	1

Number of holdings to be sampled of each poultry category (except turkey, duck and goose holdings)

Number of holdings to be sampled
All
35
42
53
60

Number of holdings per Member State	Number of holdings to be sampled
Up to 46	All
47-60	47
61-100	59
101-350	80
> 350	90

Table 2

Number of turkey, duck and goose holdings to be sampled

C. Specific requirements for detection of infections with H5/H7 subtypes of avian influenza in ducks, geese and quail

1. Blood samples for serological testing shall be taken preferably from birds which are kept outside in fields.

- 2. From each selected holding 40-50 blood samples shall be taken for serological testing.
- 3. In case commercial flocks are not present, surveillance could be carried out on backyard flocks.

D. Laboratory testing

- 1. Laboratory tests shall be carried out in accordance with the avian influenza diagnostic manual (Commission Decision 2006/437/EC) laying down the procedures for the confirmation and differential diagnosis of avian influenza (including examination of sera from ducks and geese by haemagglutination-inhibition (HI) test).
- 2. However, if laboratory tests not laid down in the avian influenza diagnostic manual nor described in the OIE Terrestrial Manual are envisaged, Member States shall provide the necessary validation data to the CRL, in parallel to submitting their programme to the Commission for approval.
- 3. All positive serological findings shall be confirmed by the National Laboratories for avian influenza by a haemagglutination-inhibition test, using designated strains supplied by the Community Reference Laboratory for Avian Influenza:
 - H5 (a) initial test using Ostrich/Denmark/72420/96 (H5N2);
 - (b) test all positives with Duck/Denmark/64650/03 (H5N7) to eliminate N2 cross reactive antibody.
 - H7 (a) initial test using Turkey/England/647/77 (H7N7);
 - (b) test all positives with African Starling/983/79 (H7N1) to eliminate N7 cross reactive antibody.

ANNEX II

Guidelines on the implementation of surveillance programmes for avian influenza in wild birds to be carried out in the Member States

A. Objectives, general requirements and criteria for surveillance

A1. Objectives

Virological surveillance for avian influenza in wild birds aim to identify the risk of introduction of AI viruses (LPAI and HPAI) to domestic poultry by:

- ensuring early detection of HPAI H5N1 by investigating increased incidence of morbidity and mortality in wild birds, in particular in selected 'higher risk' species.
- in the event that HPAI H5N1 is detected in wild birds, then surveillance of live and dead wild birds shall be enhanced to determine whether wild birds of other species can act as asymptomatic carriers or 'bridge species' (see Part E of this Annex).
- continuing a 'baseline' surveillance of different species of free living migratory birds as part of continuous monitoring of LPAI viruses. Anseriformes (water fowl) and Charadriiformes (shorebirds and gulls) shall be the main sampling targets to assess if they carry LPAI viruses of H5 and H7 subtypes (which would in any case also detect HPAI H5N1 and other HPAI, if present). 'Higher risk species' must be targeted in particular.

A2. General requirements and criteria

1. Sampling shall not extend beyond 31 December of the year of implementation of the programme.

- 2. Testing of samples shall be carried out at national laboratories for avian influenza (NL) in Member States or by other laboratories authorised by the competent authorities and under the control of the NL.
- 3. All results shall be sent to the Community Reference Laboratory for Avian Influenza (CRL) for collation. A good flow of information must be ensured. The CRL shall provide technical support and keep an enlarged stock of diagnostic reagents.
- 4. All avian influenza virus isolates of cases in wild birds shall be submitted to the CRL in accordance with Community legislation, unless a derogation according to paragraph 4 of Chapter V under Differential diagnosis in the avian influenza Diagnostic Manual laid down in Decision 2006/437/EC is granted. Viruses of H5/H7 subtype shall be submitted without delay and shall be subjected to the standard characterisation tests (nucleotide sequencing/IVPI) according to the said diagnostic manual.

B. Surveillance for avian influenza in wild birds

Design and implementation

Close cooperation with epidemiologists and ornithologists and the competent authority for nature conservation shall be ensured for designing the surveillance, assisting in species identification and optimising sampling. The design of the surveillance shall be adapted to the national situation as regards selection of species to be sampled according to species predominance and bird population sizes. Sampling must consider the seasonality of migration patterns, which may vary in different Member States. It shall take into account the behaviour of bird species as regards migratory flyways, main habitats, gregariousness and degree of mixing during migration and the results obtained from previous surveillance during 2003-2006. In addition, continuous review and feedback will be provided through the AI wild bird surveillance working group who are analysing new data as they become available.

For H5N1 HPAI, all those factors shall be considered in relation to the probability of wild bird exposure to infected poultry and wild birds in outbreak areas and the probability of contact of wild birds with domestic poultry in the poultry husbandry systems in the different Member States.

To assess those probabilities, the decision trees and tables in the opinion of EFSA (¹), which were drawn up in collaboration with the European Commission's Environment Directorate-General can provide an effective tool for Member States' local risk assessments to adapt to an evolving situation based on a close collaboration and exchange of views between Member States.

Liaisons with bird conservation/watching institutions and ringing stations shall be encouraged. Sampling, where appropriate, shall be carried out under the supervision of staff from these groups/stations, by hunters and other ornithologically skilled persons.

- 1. Passive surveillance of sick and dead wild birds shall be targeted on:
 - (a) areas where increased incidence of morbidity and mortality in wild birds occurs;
 - (b) areas close to the sea, lakes and waterways where birds were found dead; and in particular when these areas are in proximity to domestic poultry farms;
 - (c) birds belonging to identified 'higher risk' species listed in part D and other wild birds living in close proximity with them.
- 2. In addition, investigations of living and dead wild birds in areas where H5N1 cases have been detected shall ideally be targeted on birds:
 - (a) wild birds or poultry to possibly identify asymptomatic carriers;
 - (b) in areas epidemiologically linked to these cases;
 - (c) coming possibly in close contact to domestic poultry holdings (Protection zone, surveillance zone and area B), which might function as 'bridge species', in particular those that are listed in part E.
- 3. Active surveillance on living and clinically healthy and/or clinically diseased, injured or hunted (2) birds shall be targeted on:
 - (a) migratory birds belonging to the orders of Anseriformes (water fowl) and Charadriiformes (shorebirds and gulls);
 - (b) at identified areas for concentration and mixing of high number of migratory birds involving different species and in particular when these areas are in proximity to domestic poultry farms;
 - (c) a selection of higher risk species (3).

Sampling procedures

 Oropharyngeal and cloacal swabs for virological examination shall be taken from apparently healthy free living birds. If for any reason it is impractical to take cloacal swabs from live birds carefully collected fresh faeces samples may serve as an alternative. However, traceability in case of mixed sites frequented by different bird species must be ensured.

Scientific opinion on 'Migratory birds and their possible role in the spread of highly pathogenic avian influenza' (EFSA 12 May 2006).
 Hunting by respecting the requirements of Council Directive 79/409/EEC on the protection and conservation of all naturally occurring

wild birds.

 $^(^{3})$ To be provided by the European Commission's Environment Directorate-General.

- Cloacal and tracheal/oropharyngeal swabs and/or tissues (namely the brain, heart, lung, trachea, kidney and intestines) from wild birds found dead or shot shall be sampled for virus isolation and molecular detection (PCR).
- 3. Specific care has to be taken for the storage and transport of samples. Swabs must be chilled immediately on ice or with frozen gel packs and submitted to the laboratory as quickly as possible. Samples must not be frozen unless absolutely necessary. If available, swabs must be placed in antibiotic or specific virus transport medium so that they are fully immersed. Placing samples in medium for transportation must be done in addition to chilling and not as an alternative to chilling. In the absence of such medium, swabs must be returned to their casing and submitted dry. If rapid transport within 48 hours to the laboratory (in transport medium at 4° Celsius) is not guaranteed, samples shall be immediately frozen, stored and then transported on dry ice. Storage and transport of samples may be affected by a variety of factors so the method selected must be fit for purpose.
- 4. Sampling procedures shall be carried out in accordance with the avian influenza diagnostic manual (Commission Decision 2006/437/EC) laying down the procedures for the confirmation and differential diagnostic of avian influenza.

C. Laboratory testing

- 1. Laboratory tests shall be carried out in accordance with the avian influenza diagnostic manual (Commission Decision 2006/437/EC) laying down the procedures for the confirmation and differential diagnostic of avian influenza.
- 2. However, if laboratory tests not laid down in the avian influenza diagnostic manual nor described in the OIE Terrestrial Manual are envisaged, Member States shall provide the necessary validation data to the CRL, in parallel to submitting their programme to the Commission for approval.
- 3. All samples collected in the surveillance for avian influenza in wild birds shall be tested as soon as possible by molecular techniques if available and according to the diagnostic manual (Commission Decision 2006/437/EC). These tests shall only be carried out in laboratories able to guarantee quality assurance and using methods recognised by the CRL for avian influenza. In addition, methods used must have produced acceptable results in the most recent comparative ring test of national laboratories. Initial screening using M gene PCR is recommended, with rapid testing of positives for H5 (but within two weeks) and in case of a positive finding analysis of the cleavage site must be undertaken as soon as possible to determine whether or not it has a highly pathogenic avian influenza (LPAI) or a low pathogenic avian influenza (LPAI) motif. If H5 HPAI is confirmed further analysis to determine the N type must be done rapidly (even this can only provide evidence eliminating N1).
- 4. At the laboratory, pooling of up to five samples taken from the same species collected at the same site and same time may be permitted when it can be ensured that, in case of a positive finding, the individual samples can be identified and retested.
- 5. Serological surveillance shall not be applied for avian influenza investigations in wild birds because serological methods cannot distinguish between HP and LP strains and antibody findings do not allow inference in relation to the likely location where wild birds might have become infected. However, serological surveillance might be important to study in which resident or migrating bird species H5/H7 viruses are/were prevalent (or endemic). Such analysis shall only be performed by specialised laboratories using a carefully selected panel of antigens to ensure the detection of haemagglutinin specific antibodies (i.e. to eliminate the possibility of interference from N specific antibodies).

D. List of wild bird species presenting a higher risk in relation to avian influenza (*)

Common name	Scientific name		
Bewick's Swan	Cygnus columbianus		
Whooper Swan	Cygnus cygnus		
Mute Swan	Cygnus olor		
Geese			
Pink-footed Goose	Anser brachyrhynchus		
Bean Goose	Anser fabalis		
Greater White-fronted Goose (European race)	Anser albifrons albifrons		
Lesser White-fronted Goose	Anser erythropus		
Greylag Goose	Anser anser		
Barnacle Goose	Branta leucopsis		
Brent Goose	Branta bernicla		
Red-breasted Goose	Branta ruficollis		
Canada Goose	Branta canadensis		
Ducks			
Eurasian Wigeon	Anas penelope		
Common Teal	Anas crecca		
Mallard	Anas platyrhynchos		
Northern Pintail	Anas acuta		
Garganey	Anas querquedula		
Northern Shoveler	Anas clypeata		
Marbled Teal	Marmaronetta angustirostris		
Red-crested Pochard	Netta rufina		
Common Pochard	Aythya ferina		
Tufted Duck	Aythya fuligula		
Waders			
Northern Lapwing	Vanellus vanellus		
Eurasian Golden Plover	Pluvialis apricaria		
Black-tailed Godwit	Limosa limosa		
Ruff	Philomachus pugnax		
Gulls			
Black-headed Gull	Larus ridibundus		
Common Gull	Larus canus		
(*) This list is not a limitating list but is only meant to iden	tify migratory species that may pose a high risk for introduction of avian		

*) This list is not a limitating list but is only meant to identify migratory species that may pose a high risk for introduction of avian influenza into the Community based on their migratory pattern involving areas where H5N1 HPAI has either occurred in wild birds or poultry. It is based on the Scientific opinion on 'Migratory birds and their possible role in the spread of highly pathogenic avian influenza' adopted by the Animal Health and Welfare Panel of EFSA on 12 May 2006 and the work carried out by ORNIS Committee and contractors to European Commission's Environment Directorate-General. However, this list could be updated following results of further scientific studies as they become available and based on the risk assessment carried out by national authorities taking into account their specific ornithological situation.

E. List of birds living in proximity to domestic poultry (**)

Common name	Scientific name	Probability of contact with poultry			
Group 1. Species intimately associated with poultry production in Europe					
Domestic Goose	Anser anser domesticus	High			
Domestic Mallard	Anas platyrhynchos	High			
Domestic Muscovy Duck	Cairina moschata	High			
Feral Pigeon	Columba livia	High			
House Sparrow	Passer domesticus	High			
Group 2. Species which may share	farmland also used by domesticated	l poultry in north Europe			
Eurasian Golden Plover	Pluvialis apricaria	Low			
Northern Lapwing	Vanellus vanellus	Medium			
Black-headed Gull	Larus ridibundus	High			
Common Gull	Larus canus	High			
Herring Gull	Larus argentatus	Low			
Wood Pigeon	Columba palumbus	High			
Eurasian Collared Dove	Streptopelia decaocto	High			
Ring-necked Pheasant	Phasianus colchicus	High			
Larks species	Alauda & Galerida spp	Low			
Pipits		Low			
Wagtails		Medium			
Fieldfare	Turdus pilaris	Medium			
Redwing	Turdus iliacus	Medium			
Black-billed Magpie	Pica pica	High			
Eurasian Jackdaw	Corvus monedula	High			
Rook	Corvus frugilegus	Medium			
Carrion Crow	Corvus corone	Medium			
Raven	Corvus corax	Low			
Starling	Sturnus vulgaris	High			
Spotless Starling	Sturnus unicolor	High			
House Sparrow	Passer domesticus	High			
Eurasian Tree Sparrow	Passer montanus	High			
Finches		Medium			
Buntings	Miliaria, Emberiza spp	Medium			

Common name	Scientific name	Probability of contact with poultry

Group 3. Species which may share wetlands also used by domesticated water birds in Northern Europe

Egrets	Egretta spp.	Low
Herons	Ardea and other spp.	Medium
Cormorant	Phalacrocorax carbo	Medium
Storks	Ciconia spp.	Low
Mute Swan	Cygnus olor	Medium
Greylag Goose	Anser anser	Medium
Canada Goose	Branta canadensis	Low
Ducks	Anas & Aythya spp.	Low
Mallard	Anas platyrhynchos	High
Common Coot	Fulica atra	Medium
Moorhen	Gallinula chloropus	Medium

(**) This list is not an exhaustive list but is only meant to identify resident or non-migratory European bird species that live in proximity to domestic poultry (especially in NW Europe) and which have the theoretical potential to transfer HPAI H5N1 from potential asymptomatically infected wild birds ('bridge species'). It is based on the Scientific opinion on 'Migratory birds and their possible role in the spread of highly pathogenic avian influenza' adopted by the Animal Health and Welfare Panel of EFSA on 12 May 2006 and the work carried out by ORNIS Committee and contractors by European Commission's Environment Directorate-General (DG ENV). However, this list could be updated and expanded following results of further scientific studies as they become available. In particular, DG ENV has commissioned Wetland international and EURING to review, update and expand the preliminary analysis of higher risk species and sites in the light of H5N1 outbreaks in Europe in 2006, and to identify other High Risk bird species which might act as 'bridge species' between wild birds and poultry and/or humans in different parts of Europe. The results should be available by end of June 2007.

(http://ec.europa.eu/environment/nature/nature_conservation/focus_wild_birds/avian_influenza/pdf/avian_influenza_report.pdf). It should provide a much more reliable preliminary list of higher risk species and contact risk with poultry within the EU and provides for a more targeted approach.

ANNEX III

'ANNEX IV

Standard requirements for the submission of surveillance programmes for avian influenza co-financed by the Community

Identification of the programme 1.

Member State:

Disease:

Year of implementation:

Reference of this document:

Contact (name, phone, fax, e-mail):

Date sent to the Commission:

2. Description of the surveillance programme in poultry

- 2.1. Objectives, general requirements and criteria
- 2.2. Design and implementation

Table 2.2.1

Poultry holdings (a) (except ducks and geese) to be sampled

Serological investigation according to point B of Annex I to Commission Decision 2007/268/EC on holdings of broilers (only when at risk)/fattening turkeys/chicken breeders/turkey breeders/laying hens/free range laying hens/ratites/farmed feathered game (pheasants, partridges, quails...)/"backyard flocks"/others (delete as appropriate)

PLEASE USE ONE FORM PER POULTRY CATEGORY

NUTS (2) code (^b)	Total number of holdings (°)	Total number of holdings to be sampled	Number of samples per holding	Total number of tests to be performed per method	Methods of laboratory analysis
Total					

Holdings equals herds, flocks or establishments as appropriate. (a)

(b) Refers to the location of the holding of origin. In case Nuts 2 code can not be used, coordinates (long/lat) are requested.
 (c) Total number of holdings of one category of poultry in concerned NUTS 2 region.

Table 2.2.2

Duck and geese holdings to be sampled (a) according to point C of Annex I to Decision 2007/268/EC

Serological investigation

NUTS (2) code (^b)	Total number of duck and geese holdings	Total number of duck and geese holdings to be sampled	Number of samples per holding	Total number of tests to be performed per method	Methods of laboratory analysis
Total					

(a) Holdings equals herds, flocks or establishments as appropriate.

(*) Refers to the location of the holding of origin. In case Nuts 2 code can not be used, coordinates (long/lat) are requested.

2.3. Laboratory testing: description of the laboratory tests used

3. Description of the surveillance programme in wild birds:

- 3.1. Objectives, general requirements and criteria
- 3.2. Design and implementation

Table 3.2.1

Wild birds – investigation according with the programme for surveillance of avian influenza in wild birds set out in Annex II to Decision 2007/268/EC

NUTS (2) code (ª)	Wild birds to be sampled (^b)	Total number of samples to be taken for active surveillance	Total number of samples to be taken for passive surveillance
Total			

(a) Refers to the place of collection of birds/samples. In case Nuts 2 code can not be used, coordinates (long/lat) are requested.
 (b) General description of the wild birds are intended to be sampled in the framework of the active and passive surveillance.

3.3. Laboratory testing: description of the laboratory tests used

4. Description of the epidemiological situation of the disease in poultry during the last five years

- 4.1. Measures included in the programme for poultry surveillance
- 4.1.1. Designation of the central authority charged with supervising and coordinating the departments responsible for implementing the programme
- 4.1.2. System in place for the registration of holdings

4.1.3. Data on vaccination

5. Description of the epidemiological situation of the disease in wild birds during the last five years

- 5.1. Measures included in the programme for wild birds surveillance
- 5.1.1. Designation of the central authority charged with supervising and coordinating the departments responsible for implementing the programme
- 5.1.2. Description and delimitation of the geographical and administrative areas in which the programme is to be applied
- 5.1.3. Estimation of the local and/or migratory wildlife population

6. Measures in place as regards the notification of the disease

- 7. Costs
- 7.1. Detailed analysis of the costs:
- 7.1.1. Poultry
- 7.1.2. Wild birds
- 7.2. Summary of the costs
- 7.2.1. Poultry surveillance

Measures eligible for co-financing surveillance in poultry				
Methods of laboratory analysis	Number of tests to perform per method	Unitary test cost (per method)	Total cost	
Serological pre-screening (ª)				
Haemagglutination-inhibition-test (HI) for H5/H7 (^b)				
Virus isolation test				
PCR test				
Other measures to be covered	Specify activities			
Sampling				
Others				
Total				
(a) Specify the loberatory test to be used	· · · · ·		1	

(b) Specify number of tests for H5 and for H7.

7.2.2. Wild bird surveillance

Measures eligible for co-financing surveillance wild birds				
Methods of laboratory analysis	Number tests to perform per method	Unitary test cost (per method)	Total cost	
Serological pre-screening				
Haemagglutination-inhibition-test (HI) for H5/H7				
Virus isolation test				
PCR test				
Other measures to be covered	Specify activities			
Sampling				
Others				
Total				